Surname	Centre Number	Candidate Number
First name(s)		0



GCSE

3410U20-1



FRIDAY, 27 MAY 2022 - MORNING

CHEMISTRY – Unit 2:

Chemical Bonding, Application of Chemical Reactions and Organic Chemistry

FOUNDATION TIER

1 hour 45 minutes

For Examiner's use only		
Question	Maximum Mark	Mark Awarded
1.	9	
2.	7	
3.	7	
4.	13	
5.	10	
6.	6	
7.	8	
8.	6	
9.	7	
10.	7	
Total	80	

ADDITIONAL MATERIALS

In addition to this examination paper you will need a calculator and a ruler.

INSTRUCTIONS TO CANDIDATES

Use black ink or black ball-point pen. Do not use gel pen or correction fluid.

You may use a pencil for graphs and diagrams only.

Write your name, centre number and candidate number in the spaces at the top of this page. Answer **all** questions.

Write your answers in the spaces provided in this booklet. If you run out of space, use the additional page(s) at the back of the booklet, taking care to number the question(s) correctly.

INFORMATION FOR CANDIDATES

The number of marks is given in brackets at the end of each question or part-question.

Question **6** is a quality of extended response (QER) question where your writing skills will be assessed.

The Periodic Table is printed on the back cover of this paper and the formulae for some common ions on the inside of the back cover.



[4]

Answer all questions.


1. Selwyn carried out an experiment to prepare copper(II) chloride crystals.


The diagrams show the stages of the experiment he carried out.


(a) Choose apparatus from the box to label the diagrams.

evaporating basin conical flask filter funnel electronic balance filter paper beaker test tube

bubbles of gas









© WJEC CBAC Ltd.

PMT

	3410U201	0.3

		Examin
(b)	In Stage 1 Selwyn added copper(II) carbonate until all the dilute hydrochloric acid was used up.	only
	Tick $(\mspace{1mu})$ the box next to the statement which best describes what Selwyn would see when all the acid had been used up.	1]
	bubbling increases	
	bubbling stops	
	bubbling decreases	
(c)	The gas formed in Stage 1 turns limewater milky. <u>Underline</u> the name of this gas. [1]
	oxygen hydrogen carbon dioxide nitrogen	
(d)	Choose words from the box to complete the following sentences. [2	2]
	evaporation filtration distillation neutralisation	
	The process used to remove the unreacted copper(II) carbonate in Stage 2 is called	
	The process used to remove water in Stage 3 is called	
(e)	The container holding the crystals has a mass of 29.8 g. Using the information given in Stage 4 , calculate the mass of the crystals formed.	1]
	Mass of crystals formed =	9

(3410U20-1) © WJEC CBAC Ltd.

Turn over.

2.	(a)	Crude oil is formed from simple marine organisms.	Examiner only
		Tick (\(\strict{\strict} \)) the box next to the length of time it takes for the organisms to turn into crude oil.	[1]
		hundreds of years	ניז
		thousands of years	
		millions of years	
	(b)	Crude oil can be separated into simpler mixtures called fractions.	
		Tick (\(\strict{\strain} \)) the box next to the method used to separate crude oil into fractions.	[1]
		fractional distillation	
		filtration	
		cracking	
		polymerisation	



PMT

(c) The table shows properties of some fractions that are obtained from crude oil.

Fraction	Size of molecules (chain length)	Viscosity	Ease of ignition	Amount of smoke formed
petrol	C ₅ -C ₁₀	very runny	very easy	no smoke
naphtha	C ₈ -C ₁₂	fairly runny	quite easy	little smoke
kerosene	C ₁₀ -C ₁₆	thick	quite hard	quite a lot of smoke
diesel oil	C ₁₄ -C ₂₀	very thick	very hard	very smoky

Use only the information from the table to answer the following questions.

(i)	Name the fraction which is the easiest to pour.	[1]
(ii)	Name the fraction which is the hardest to burn.	[1]
(iii)	Name the fraction with the smallest range of chain lengths.	[1]

(iv) Name the fraction which burns with the cleanest flame. Give the reason for your choice. [2]

Fraction

Reagon	
INCASOII	

7



[2]

PMT

3. (a) Complete the table.

Name	Molecular formula	Structural formula
ethane	C ₂ H ₆	H H
propane	C ₃ H ₈	
butane		H H H H

(b) The box contains four polymers **A**, **B**, **C** and **D**.

Give the **letter** of the polymer formed during the polymerisation of ethene. [1]



PMT

(c) The equation shows the reaction taking place when testing for a C = C bond.

bromine water potassium dichromate universal indicator barium chloride

Choose the name of reagent **X** from the box above.

.....[1]

(d) The equation shows the fermentation of glucose.

$$C_6H_{12}O_6$$
 yeast $2C_2H_5OH + 2CO_2$

(i) <u>Underline</u> the chemical name for C_2H_5OH .

[1]

sugar biofuel ethanol alcohol

(ii) Calculate the relative molecular mass, $M_{\rm r}$, of ${\rm C_2H_5OH}$.

[2]

$$A_r(H) = 1$$
 $A_r(O) = 16$ $A_r(C) = 12$

*M*_r =

7


BLANK PAGE

PLEASE DO NOT WRITE ON THIS PAGE



PMT

4. (a) The diagram shows the apparatus a teacher used to demonstrate what happens when an electric current is passed through molten lead bromide.



(i) Lead bromide contains the ions Pb2+ and Br -.

<u>Underline</u> the correct formula of lead bromide.

[1]

(ii) Give the state (solid, liquid or gas) of the lead bromide during the process. [1]

(iii) Name the orange gas formed at the anode. [1]


(iv) During the process each lead ion, Pb²⁺, gains two electrons forming a lead atom at the cathode. [1]

Tick (\mathcal{I}) the box next to the equation for the reaction at the cathode.

$$Pb^{2+} + 2e^{-} \longrightarrow Pb$$



(b) Iron is extracted from its ore in the blast furnace. The diagram shows the materials which enter and leave the furnace.





PMT

(i) <u>Underline</u> the correct word(s) in the brackets to complete each sentence.

[3]

The furnace is heated by burning (coke/iron/iron ore).

Hot air provides (waste gases/oxygen/slag) for burning to take place.


Impurities are removed by adding (steel/hot air/limestone) to the furnace.

(ii) The word equation shows the main reaction taking place in the blast furnace.

iron(III) oxide + carbon monoxide → iron + carbon dioxide

Give the **letter** next to the products which correctly balance the symbol equation for this reaction. [1]

+ 3CO₂



Letter



(c) Chemists have designed a wide variety of alloys for different uses.Some alloys contain iron and carbon only, whereas others contain additional metals.The table shows the composition and properties of some alloys.

Alloy	Composition	Properties
mild steel	iron plus 0.15-0.30% carbon	malleable (easy to bend), ductile (easy to pull into wire) and soft (easy to scratch)
high carbon steel	iron plus 0.70-1.50% carbon	strong, brittle and hard
cast iron	iron plus 2.00-5.00% carbon	very strong, very brittle and very hard

Use only the information in the table above to answer parts (i)-(iii).


(1)	warrie trie alloy triat contains trie leas	a amount of carbon.	ניו
(ii)	<u>Underline</u> the effect of increasing the	percentage (%) of carbon in these alloys.	[1]
	strength decreases	hardness decreases	
	softness increases	brittleness increases	
(iii)	Give the property of mild steel which	makes it useful for making car bodies.	[1]



PMT


(iv) Diagrams **A**, **B** and **C** are models showing the arrangement of atoms in pure iron and in two alloys, **but not necessarily in that order**.

Choose the **letter** of the model which best represents cast iron. Give the reason for your choice. [2]



Α

В



Letter

Reason

341

13




© WJEC CBAC Ltd. (3410U20-1)

Turn over.

5. Joe, Alex and Megan were asked to investigate the temperature rise when four metals were added to excess copper(II) sulfate solution. 0.5 g of each metal was added to separate 50 cm³ samples of copper(II) sulfate solution.

The temperature rise for each reaction was measured.



Their results are shown in the table below.


Metal	Temperature rise (°C)
magnesium	47
calcium	54
zinc	38
iron	25



© WJEC CBAC Ltd.

PMT

(a) (i) Draw a bar chart on the grid to show the results. One bar has been drawn for you.



(ii) Put the **four** metals in order of reactivity. [1]

Most reactive

.....

Least reactive

15

© WJEC CBAC Ltd. (3410U20-1) Turn over.

3410U201

(iii)	The temperature rise for each metal was lower than expected. The students we asked to suggest how the apparatus could be improved so that the temperature rise recorded for each metal was closer to the expected value.	
	Joe suggested wrapping the polystyrene cup in cotton wool.	
	Megan suggested putting a lid on the polystyrene cup.	
	Alex suggested using a cup made from copper.	
	Choose which student's suggestion would not result in the temperature rises being closer to the expected value. Explain your choice.	[2]
	Student	
	Reason	
•••••		
(iv)	Complete the equation for the reaction between magnesium and copper(II) sulfate solution.	[2]
	Mg + CuSO ₄ +	




© WJEC CBAC Ltd.

(b)	The students repeated the experiment using chromium and recorded a temperature rise of 30 °C.		Examinei only
	What does the temperature rise of 30 °C tell you about the reactivity of chromium compared to the four metals in part (a)?	[1]	
(c)	The energy given out by the reaction can be calculated using the formula below.		
	energy given out (J) = volume of solution (cm 3) \times 4.2 \times temperature rise (°C)		
	Calculate the energy given out during the displacement reaction between iron and copper(II) sulfate solution.	[2]	
	Energy given out =	J	
			10
		- 1	



Turn over.

6. The picture below shows the universal indicator pH colour chart and the position of some substances on the pH scale.





© WJEC CBAC Ltd.

Describe what is meant by the pH scale and what it tells us about the substances shown in the picture. [6 QER	Exa



© WJEC CBAC Ltd. (3410U20-1) Turn over.

7. (a) Skiers and mountaineers use hand warmers in very cold weather conditions. Hand warmers are small packets that produce heat when needed. Some ski clothing is specially designed to hold a hand warmer.





Table 1 describes how three types of hand warmers work.

A	B	C
Air-activated	Battery powered	Supersaturated solution
(disposable)	(reusable)	(reusable)
The packaging seal is broken allowing air to reach the chemicals causing a chemical reaction to occur. This type of hand warmer can only be used once.	A metal coil heats up when the device is switched on. The device needs recharging.	A metal button on the packaging is pressed. This causes crystals to form in the solution. The hand warmer can be reactivated by placing it in boiling water.

Table 1



© WJEC CBAC Ltd.

Table 2 shows some information about each type of hand warmer.

Type of hand warmer	Cost	Time to warm			re after beir ced in a fre		I
	(£)	(£) up after activation	after 15 mins	after 30 mins	after 60 mins	after 90 mins	after 120 mins
Α	1	less than 1 min	39	39	38	38	37
В	80	less than 1 min	32	29	28	27	26
С	24	less than 1 min	42	34	27	18	8

Table 2

(i)	Tick (✓) the box next to the correct statement.	[1]
	Gloves need to be worn when using hand warmers	
	Boiling water is used to recharge battery powered hand warmers	
	Some chemical reactions give out heat energy	
	All hand warmers are reusable	
(ii)	Give two reasons why hand warmer A is the most popular choice.	[2]
	Reason 1	
	Reason 2	
•••••		



© WJEC CBAC Ltd. (3410U20-1) Turn over.

(b) An air-activated hand warmer contains several chemicals mixed together. One of these chemicals is iron.

When the pack is opened, a reaction takes place between iron and oxygen in the air, causing an increase in mass.

A student investigated the increase in mass of the opened pack over several hours. The results are shown below.

Time (hours)	Increase in mass (g)
0.0	0.0
0.5	1.4
1.0	2.6
1.5	3.7
2.0	4.4
2.5	4.7
3.0	4.8
3.5	4.8
4.0	4.8



Examiner only Plot the mass increase of the pack against time on the grid and draw a suitable line. Two plots have been done for you. [3] 5 4 3 Increase in mass (g) 2 Time (hours) (ii) Tick (\mathcal{I}) the box next to the time it takes for the reaction to finish. [1] 2 hours 3 hours 4 hours 5 hours (iii) Tick $(\mbox{\em 4})$ the box next to the statement that best explains the shape of the graph. [1] Iron reacts with oxygen forming iron oxide until all the oxygen is used up Heat formed expands the iron Iron oxide loses oxygen, forming iron Iron reacts with oxygen forming iron oxide until all the iron is used up 8




© WJEC CBAC Ltd. (3410U20-1) Turn over.

(a) The table shows the electronic structure of the elements present in potassium fluoride.

Examiner only

Element	Electronic structure
potassium	2,8,8,1
fluorine	2,7

The diagram shows the electron transfer that occurs when potassium reacts with fluorine to form potassium fluoride. The ● and × symbols are outer shell electrons.



A student was asked to draw a diagram showing the electronic structures and charges on the ions formed. There are **two** mistakes in the student's answer.


Student's answer (2,8,8) (2,8,8)

- (i) Circle the **two** mistakes in the student's answer. [2]
- (ii) Name the type of bonding found in potassium fluoride. [1]



8.


(iii) The diagrams show four different structures. Give the **letter** of the structure most likely to represent potassium fluoride. [1]






В

Α



C




Letter

(b) The diagram shows the electrons in the outer shell of an atom of fluorine.



Complete the diagram to show the outer shell electrons in a molecule of fluorine. [2]




6

9. (a) The diagram outlines the manufacture of ammonia by the Haber process.



[2]



- (i) Name the raw material **X**. [1]
- (ii) The pressure used in the Haber process is 200 atm. State why a higher pressure is **not** used. [1]
- (iii) At 450 °C, the reaction is very slow. Iron is used in the process to speed up the reaction. Give the name for a substance used to speed up a chemical reaction. [1]
- (iv) The reaction between nitrogen and hydrogen is represented by the equation below.

$$N_2$$
 + $3H_2$ \Longrightarrow $2NH_3$

Complete the equation below using the key:

nitrogen gas, N₂





26

(b) One of the main uses of ammonia is in the manufacture of fertilisers.

The table shows the results obtained when tests were carried out on three different fertilisers ${\bf A},\,{\bf B}$ and ${\bf C}.$

Fertiliser	Test for positive ion	Test for negative ion
A	On adding sodium hydroxide solution and warming, a pungent smelling gas is formed which turns red litmus blue	On adding barium chloride solution a white precipitate forms
В	Lilac flame test	On adding silver nitrate solution a white precipitate forms
С	On adding sodium hydroxide solution and warming, a pungent smelling gas is formed which turns red litmus blue	On adding silver nitrate solution a white precipitate forms

Give the letter of the fertiliser which is ammonium sulfate.	[1]
Letter	

(c) Ammonia reacts with chlorine to form nitrogen and hydrogen chloride.

Complete the balancing of the equation for this reaction.

2NH ₃	+	3Cl ₂	 N_2	+	HCI

7

[1]

10.	(a)	The	ist below shows part of the reactivity series.	Examine only
			sodium	
			aluminium	
			(carbon)	
			tin	
			copper	
			silver	
		(i)	Tin is extracted from its ore by heating with carbon. Aluminium is extracted from its ore using a different method. Give the name of the method used to extract aluminium.	
		(ii)	The equation shows the extraction of tin from tin oxide using carbon.	
			$SnO_2 + C \longrightarrow Sn + CO_2$	
			Tick (✓) the box next to the correct statement. [1]	
			Carbon is reduced	
			Tin is oxidised	
			Tin oxide is reduced	
			Carbon dioxide is oxidised	
		(iii)	When aluminium and copper(II) oxide are heated together, aluminium oxide and copper are formed.	
			Complete and balance the equation for this reaction. [3]	
		2A	l + 3 + Cu	
			copper(II) oxide aluminium oxide	



© WJEC CBAC Ltd.

(b) A teacher wanted to find out the position of four metals A, B, C and D in the series. She heated each metal in turn with oxides of the other three. The results of follows. A reduced the oxide of C B reduced the oxide of A B reduced the oxide of B Place the metals in order of reactivity. Most reactive	
A reduced the oxide of C B reduced the oxide of A B reduced the oxide of C D reduced the oxide of B Place the metals in order of reactivity. Most reactive	
B reduced the oxide of C B reduced the oxide of C D reduced the oxide of B Place the metals in order of reactivity. Most reactive	[2]
B reduced the oxide of C D reduced the oxide of B Place the metals in order of reactivity. Most reactive	[2]
D reduced the oxide of B Place the metals in order of reactivity. Most reactive	[2]
Place the metals in order of reactivity. Most reactive	[2]
Most reactive	[2]
Least reactive	
END OF PAPER	



Turn over.

Question number	Additional page, if required. Write the question number(s) in the left-hand margin.						
	Time the question named (o) in the left hand margin.	only					
		[
		[
		1					



FORMULAE FOR SOME COMMON IONS

POSITIV	E IONS	NEGATIVE IONS		
Name	Formula	Name	Formula	
aluminium	Al ³⁺	bromide	Br ⁻	
ammonium	$\mathrm{NH_4}^+$	carbonate	CO ₃ ²⁻	
barium	Ba ²⁺	chloride	CI ⁻	
calcium	Ca ²⁺	fluoride	F ⁻	
copper(II)	Cu ²⁺	hydroxide	OH ⁻	
hydrogen	H⁺	iodide	17	
iron(II)	Fe ²⁺	nitrate	NO ₃ -	
iron(III)	Fe ³⁺	oxide	O ²⁻	
lithium	Li⁺	sulfate	0 ²⁻ SO ₄ ²⁻	
magnesium	Mg ²⁺		·	
nickel	Ni ²⁺			
potassium	K ⁺			
silver	Ag^{+}			
sodium	Na ⁺			
zinc	Zn ²⁺			



Krypton 36 Xe Xenon 54

222 **Rn** Radon 86

20 **Ne** Ne on

19 Fluorine 9

16 O Oxygen 8

14 **N** Nitrogen

12 C Carbon 6

11 Boron 5

40 Argon

35.5 CI

S 33

₽ ₽

Si 58

²⁴

THE PERIODIC TABLE Group


9

S

က

Hydrogen

-				,		_	
Chlorine 17	80 Br	Bromine 35	127	lodine 53	210 At	Astatine 85	
Sulfur 16	79 Se	Selenium 34	128 Te	Tellurium 52	210 Po	Polonium 84	
Phosphorus 15	75 As	Arsenic 33	122 Sb	Antimony 51	209 Bi	Bismuth 83	
Silicon 14	73 Ge	Germanium 32	119 Sn	Tin 50	207 Pb	Lead 82	
Aluminium 13	70 Ga	Gallium 31	115 –	Indium 49	204 TI	Thallium 81	
	65 Zn	Zinc 30	112 Cd	Cadmium 48	201 Hg	Mercury 80	
	63.5 Cu	Copper 29	108 Ag	Silver 47	197 Au	Gold 79	
	59 Z	Nickel 28	106 Pd	Palladium 46	195 Pt	Platinum 78	
	59 Co	Cobalt 27	103 Rh	Rhodium 45	192 Ir	Iridium 77	
	56 F e	Iron 26	101 Ru	Ruthenium 44	190 Os	Osmium 76	Key
	55 Mn	Manganese 25	99 Tc	Technetium 43	186 Re	Rhenium 75	
	⁵² Cr	Chromium 24	96 Mo	Molybdenum 42	184 W	Tungsten 74	
	51	Vanadium 23	93 Nb	Niobium 41	¹⁸¹ Ta	Tantalum 73	
	48 Ti	Titanium 22	91 Zr	Zirconium 40	179 Hf	Hafnium 72	
	45 Sc	Scandium 21	% % >	Yttrium 39	139 La	Lanthanum 57	227 Ac Actinium 89
Magnesium 12	40 Ca	Calcium 20	88 Sr	Strontium 38	137 Ba	Barium 56	226 Ra Radium 88
Sodium 11	₩ 30	Potassium 19	% Rb	Rubidium 37	133 Cs	Caesium 55	223 Fr Francium 87



© WJEC CBAC Ltd.

9 **Be** Beryllium

 \overline{Mg}^{24}